

Circadian Rhythms in the Pathogenesis and Treatment of Fatty Liver Disease

Anand R. Saran¹Shravan Dave¹Amir Zarrinpar^{1,2,3,4}

From the ¹Division of Gastroenterology, University of California, San Diego, La Jolla, California; ²Veterans Affairs Health Sciences San Diego, La Jolla, California; ³Institute of Diabetes and Metabolic Health, University of California, San Diego, La Jolla, California; and ⁴Center for Microbiome Innovation, University of California, San Diego, La Jolla, California

Circadian clock proteins are endogenous timing mechanisms that control the transcription of hundreds of genes. Their integral role in coordinating metabolism has led to their scrutiny in a number of diseases, including nonalcoholic fatty liver disease (NAFLD). Discoordination between central and peripheral circadian rhythms is a core feature of nearly every genetic, dietary, or environmental model of metabolic syndrome and NAFLD. Restricting feeding to a defined daily interval (time-restricted feeding) can synchronize the central and peripheral circadian rhythms, which in turn can prevent or even treat the metabolic syndrome and hepatic steatosis. Importantly, a number of proteins currently under study as drug targets in NAFLD (sterol regulatory element-binding protein [SREBP], acetyl-CoA carboxylase [ACC], peroxisome proliferator-activator receptors [PPARs], and incretins) are modulated by circadian proteins. Thus, the clock can be used to maximize the benefits and minimize the adverse effects of pharmaceutical agents for NAFLD. The circadian clock itself has the potential for use as a target for the treatment of NAFLD.

Keywords: Dyssynchrony; Gut Microbiome; Steatohepatitis; Lipogenesis; Bile Acids.

The 2017 Nobel Prize in Medicine signaled a recognition of the fundamental role that circadian clock proteins play in physiology and disease. It was awarded to Jeffrey C. Hall, Michael Rosbash, and Michael W. Young for their discovery of molecular mechanisms controlling circadian rhythms in *Drosophila*. Clock proteins are ubiquitous endogenous timing mechanisms that help maintain energy homeostasis by coordinating cellular processes within and between organ systems. They do so by regulating the transcription of hundreds of genes, including themselves. Their integral role in coordinating metabolism has led to closer scrutiny of their contribution to a number of diseases, including nonalcoholic fatty liver disease (NAFLD).

Although the term *circadian rhythms* usually conjures thoughts of jet lag and sleep deprivation, physiologically, they are core cellular processes that affect every organ system. The pervasiveness of clock proteins, which are found in diverse organisms from cyanobacteria to mammals, shows their evolutionary importance to life. Development of a molecular program to anticipate and prepare for periodic, predictable changes gives a tremendous adaptive advantage. Organisms have developed specialized sensory cells that allow them to adjust their own rhythmicity to external cues (called *zeitgebers*—German for “time giver”). The daily light-dark cycle on Earth governs rhythmic changes in the behavior and physiology of most species, including humans. Almost all organisms consolidate their biological processes (e.g., sleeping and feeding) to particular times of day. To acknowledge that the periodicity of these oscillations stays

Abbreviations used in this paper: ACC, acetyl-CoA carboxylase; AKT (or PKB), v-Akt murine thymoma viral oncogene homolog 1 or (protein kinase B); AMP, adenosine monophosphate; AMPK, adenosine monophosphate-activated protein kinase; BHLH, basic helix-loop-helix family member; BMAL1, brain and muscle ARNTL-like protein 1; cAMP, cyclic adenosine monophosphate; CK1, casein kinase 1; CLOCK, circadian locomotor output cycles kaput; CREB, cAMP response element-binding protein; CREBH, cAMP response element binding protein hepatocyte specific; CRY, cryptochrome; CYP7A1, cholesterol 7 alpha-hydroxylase; DBP, D-box binding protein; DEC, (or BHLH), basic helix-loop-helix family member; DIO, diet-induced obesity; ER, endoplasmic reticulum; FBXL3, F-box/LRR-repeat protein 3; FGF, fibroblast growth factor 15; FXR, farnesoid X receptor; GLP-1, glucagon-like peptide; GLUT2, glucose transporter 2; GYS2, glycogen synthase 2; HFD, high-fat diet; IRE1 α , inositol-requiring 1 α ; KO, knockout; mTOR, mammalian target of rapamycin; NAD⁺, nicotinamide adenine dinucleotide; NAFLD, nonalcoholic fatty liver disease; NFIL3 (or E4BP4), nuclear factor, interleukin 3 regulated; OR, odds ratio; OSA, obstructive sleep apnea; PER, period; PGC1 α , peroxisome proliferator-activated receptor γ coactivator 1 α ; PPAR, peroxisome proliferator-activated receptor; ROR, retinoic acid receptor-related orphan receptors; RORE, retinoic acid receptor-related-responsive element; SCFA, short-chain fatty acid; SCN, suprachiasmatic nucleus; SIRT1, sirtuin 1; SREBP, sterol regulatory element-binding proteins; T2D, type 2 diabetes; TLR, toll-like receptor; TRF, time-restricted feeding; ULK1, unc-51 like autophagy activating kinase 1; UPR, unfolded protein response.



Most current article

© 2020 by the AGA Institute
0016-5085/\$36.00

<https://doi.org/10.1053/j.gastro.2020.01.050>

approximately 24 hours, Franz Halberg, a pioneer in the field of biological rhythms, introduced the term *circadian rhythms* (from Latin, *circa* meaning “about” and *diem* meaning “a day”) in 1959.

In mammals, the circadian pacemaker resides in hypothalamic structures called the *suprachiasmatic nuclei* (SCN), and it is entrained by specialized retinal ganglion cells that measure ambient light (Figure 1). However, circadian clock proteins are not restricted to the SCN. Explants from lung, liver, and other organs exhibit sustained oscillations in vitro.^{1,2} Thus, endogenous functional clocks are also active in peripheral tissues.³ Temporal signals, such as timing of food intake, can reset the hepatic clock without affecting the SCN central clock rhythms. Consequently, different entrainment signals can activate peripheral and central circadian clocks separately.^{4,5} However, this does not undermine the importance of the SCN in coordinating physiological processes across multiple tissues.

In this review, we will describe the molecular organization of the circadian clock proteins, the relationship between circadian clock and metabolic master regulators, and clock regulation of metabolic processes important to the pathophysiology of NAFLD. Environmental cues that entrain peripheral circadian rhythms will be also discussed in detail. The effects of circadian dyssynchrony on host metabolism will be summarized, along with recent preclinical and clinical studies showing that maintaining peripheral circadian rhythms by consolidating feeding within a narrow time frame (ie, time-restricted feeding [TRF]) can restore metabolic health. Over the last 5 years, there have been many excellent reviews of preclinical data regarding the role of the circadian clock in hepatic metabolic processes specifically^{6–11} and in metabolic syndrome in general.^{12–16} Our review builds on this strong body of work by focusing on key regulatory pathways that play an important role in the treatment of steatohepatitis and fibrosis, thus showing the vital role of the circadian machinery in helping our understanding of NAFLD.

Molecular Organization of the Clock

The circadian clock in mammals, expressed in nearly every cell, is comprised of a series of transcription-translation feedback loops (Figure 2). These include circadian locomotor output cycles kaput (CLOCK) and brain and muscle ARNT-like 1 (BMAL1) as transcriptional activators. These are opposed by period (PER) and cryptochrome (CRY) as transcriptional repressors (Figure 3A). CLOCK and BMAL1 are two central components that make up the positive (or activation) limb of the molecular clock. The CLOCK:BMAL1 heterodimer then translocates into the nucleus to initiate transcription at the E-box (5'-CACGTG-3'), a specific DNA binding sequence on the promoters of its target genes.¹⁷ The main downstream targets of CLOCK:BMAL1 include their own repressors, period (PER1, PER2, and PER3) and cryptochrome (CRY1, CRY2), in addition to more than 300 other genes.¹⁸ Over time, PERs and CRYs accumulate in the cytoplasm, where they are further regulated by casein kinase 1 (CK1) ϵ , CK1 δ , and F-box/LRR-repeat protein 3 (FBXL3). CK1 ϵ and CK1 δ phosphorylate


PER for degradation; FBXL3 promotes the degradation of CRY.^{19,20} However, if CK1 ϵ phosphorylates PER after it is bound to CRY, the 3-protein complex migrates into the nucleus and inhibits the CLOCK:BMAL1 heterodimer.²¹ By inhibiting their own activators, PERs and CRYs repress the transcription of their own genes. Posttranslational modification of PERs and CRYs leads to their degradation, which then initiates a new circadian cycle with increased binding of CLOCK:BMAL1 to the now open E-box binding sites.

In a second regulatory loop, CLOCK:BMAL1 drives the transcription of the nuclear hormone receptors retinoic acid receptor-related orphan receptor (*Ror*) α and *Ror* γ and *Rev-Erb* α and *Rev-Erb* β by binding E-box elements on their promoters (Figure 3B).^{22–24} Both of these receptors can bind the ROR-responsive element (RORE) promoter DNA binding sequence (5'-AGGTCA-3'), but with opposite effects: REV-ERB α/β suppresses expression at the site, whereas ROR α/γ promotes expression.^{25,26} Together, REV-ERBs and RORs generate large-scale cyclic fluctuations in the transcription of hundreds of clock-controlled genes, including regulation of the transcription of *Bmal1*.²⁷

Recent studies have revealed an additional autonomous feedback loop involving basic helix-loop-helix family member e40 (BHLHE40; more commonly, DEC1) and BHLHE41 (more commonly, DEC2), which also suppress CLOCK:BMAL1 activity (Figure 3C).^{28,29} CLOCK:BMAL1 induces transcription of *Dec1* and *Dec2* by binding E-box elements on their promoters. Unlike PERs and CRYs, which act directly on the CLOCK:BMAL1 heterodimer, DEC2s compete with CLOCK:BMAL1 for the E-box, thus suppressing all E-box genes, including transcription of their own genes and those of PERs and CRYs.

Other circadian clock-controlled genes bring additional refinement to this complex interplay of feedback loops. CLOCK:BMAL1 modulates the expression of hundreds of other genes by promoting the transcription of D-box binding protein (DBP) by binding E-box elements on its promoter (Figure 3D).^{11,30} DBP rhythmically activates transcription of genes that have D-box elements in their promoter regions. Nuclear factor interleukin 3 (NFIL3, also E4BP4), which is regulated by a RORE promoter region (and thus is susceptible to activation by RORs and suppression by REV-ERBs), represses DBP-dependent transactivation by competing for the D-box. Computational genomic analysis of D-box elements has identified nearly 1500 regions that can be recognized by DBP and NFIL3.³¹

The transcription-translation feedback loops generate large-scale rhythms in the activities and levels of downstream clock-controlled genes.³² Because these processes are temporally controlled, it is important to regulate the abundance and activity of each component to ensure that it is in the right place at the right time. This is achieved by posttranslational modifications of clock components, such as phosphorylation, acetylation, SUMOylation (Small Ubiquitin-like Modifier proteins; [SUMO can modify the function of other proteins by attaching/detaching from them]), O-GlcNAcylation (O-Linked β -N-acetylglucosamine; [O-GlcNAc is an intracellular carbohydrate that can modify protein activity]), and ubiquitylation. Gene



expression studies show that approximately 10%–40% of the rodent genome exhibits a 24-hour rhythm in a highly tissue-specific manner.³³ However, humans are diurnal animals that feed during the day, whereas most rodents are nocturnal animals that feed primarily at night. The recent publication of a transcriptome atlas of a baboon, a diurnal primate, shows that the aforementioned percentages from rodent studies do not convey the full picture.³ Of the approximately 11,000 genes that are commonly expressed in all tissues in the baboon, 96.6% have 24-hour rhythmicity in at least 1 tissue. Greater than 80% of the 18,000 protein-coding baboon genes were circadian. More importantly, more than 80% of the genes encoding proteins targeted by pharmaceuticals are transcribed cyclically. This implies that most physiological processes are under circadian control, and thus, the disruption of circadian rhythms can result in widespread downstream pathology.

The Circadian Clock Regulates Metabolic Processes

Hepatic metabolic processes, including glucose, lipid, and cholesterol/bile acid metabolism, are highly dynamic, influenced by feeding/fasting and circadian rhythms. Data supporting these relationships come from several lines of research, including metabolic phenotyping data from circadian clock genetically modified mouse lines, molecular biology studies showing direct interactions between regulators of nutrient homeostasis and circadian clock proteins, and the relationship of feeding/fasting cycles and metabolic gene expression. In this section, we discuss the relationship between circadian clock machinery and various aspects of metabolism that are dysfunctional in patients with NAFLD.

Circadian Clock and Regulators of Nutrient Homeostasis and Autophagy

The liver contains several master metabolic regulators that sense feeding/fasting states and control the expression of hundreds of downstream metabolic targets. These include adenosine monophosphate (AMP)-activated protein kinase (AMPK), cyclic AMP (cAMP) response element-binding

protein (CREB), v-Akt murine thymoma viral oncogene homolog 1 (AKT/PKB), and sirtuin 1 (SIRT1). All have cyclic activity mainly driven by feeding behavior and can regulate—and are regulated by—circadian clock proteins.

AMPK, a fasting-sensitive protein kinase that regulates cellular energy homeostasis, can modulate, and is modulated by, the circadian clock machinery. AMPK can affect the length of circadian cycling by phosphorylating CRY and targeting it for degradation.³⁴ Moreover, ROR α activation can modulate lipid metabolism and prevent hepatic steatosis by activating AMPK, which controls key lipid regulators (discussed later in the article).³⁵ Feeding, on the other hand, activates CREB, which up-regulates *Per1* transcription.³⁶ This relationship connects G protein-coupled receptors, which use cAMP as a second messenger for transcription of circadian clock genes, with the circadian clock.³⁷ Several metabolic hormones, including glucagon and epinephrine, both of which activate hepatic gluconeogenesis, can influence circadian clock machinery through CREB activation.^{38–40} In addition, CRYs can independently suppress CREB, thereby not only affecting hepatic gluconeogenesis but also modulating *Per* transcription.³⁹ Finally, AKT, a protein kinase activated by feeding that regulates glucose metabolism, among other cellular processes (eg, cell proliferation and apoptosis), can modulate the circadian clock machinery by phosphorylating BMAL1 and CLOCK in the cytosol and preventing their translocation across the nuclear membrane.⁴¹

Sensing the nutrient status is a critical regulation step in the maintenance of homeostasis. Metabolic stressors such as starvation induce autophagy to use glycogen and lipid components for generating fuel.⁴² Circadian induction of autophagy during low nutrient states (eg, inactive state) allows efficient coordination of these cellular processes. AMPK, under glucose starvation conditions, phosphorylates and activates the autophagy-initiating kinase ULK1 (Unc-51 like autophagy activating kinase 1), thus promoting autophagy. However, under nutrient sufficiency, mammalian target of rapamycin (mTOR), another metabolic master regulator, phosphorylates ULK1 and disrupts its interaction with AMPK, thus inhibiting autophagy.⁴³ Interestingly, liver-specific *Bmal1*-null mice display dampened rhythms of autophagy gene expression, along with increased mTOR activation.⁴⁴ Thus, autophagy has

Figure 1. Central and peripheral circadian rhythms. Retinal ganglion cells detect ambient light and entrain the SCN. Their projections travel through the retinohypothalamic tract, although indirect signals are also received from the thalamus. The SCN contains the central circadian clock, which can modulate other oscillatory networks, including sleep/wake, temperature, and feeding/fasting homeostasis, through the hypothalamus. In addition, the SCN regulates the circadian release of hypothalamic-releasing hormones (eg, corticotropin-releasing hormone) and, thus, pituitary hormones (eg, adrenocorticotrophic hormone). Hormones released from the pituitary gland are one way that the central clock can regulate the peripheral organs. The SCN also regulates the peripheral clock through poorly understood neurohumoral mechanisms. SCN can drive feeding/fasting rhythms through the hypothalamus. Diet and feeding/fasting rhythms modulate cyclic fluctuations in the gut microbiome, which then create cyclic changes in secondary metabolite production, such as SCFAs and bile acids. Secondary metabolites and/or bacterial products are necessary for the maintenance of peripheral circadian rhythms. Feeding/fasting rhythms are powerful drivers of the hepatic circadian clock. The rhythmic expression of genes in the liver generates rhythmicity in the transcripts and activity of genes involved in critical metabolic processes such as glucose, lipid, and bile acid metabolism, nutrient homeostasis, autophagy, and ER stress. Metabolic syndrome occurs in the setting of central and peripheral circadian rhythm dyssynchrony.




Figure 2. Overview of circadian clock machinery. Humans/primates are diurnal animals, and their feeding/active period occurs during the light hours. However, mice (and most rodents) are nocturnal, with the feeding/active period occurring during the dark hours. The cellular molecular clock is a series of transcription-translation feedback loops that include the transcriptional activators (CLOCK and BMAL1; the positive limb) and repressors (PER and CRY; the negative limb). The PER:CRY complex repress their own activators of CLOCK:BMAL1. CLOCK:BMAL1 also promotes REV-ERBs and RORs, which add another layer of regulation over the circadian clock and regulate hundreds of additional genes.

multiple layers of regulation from metabolic regulators and the circadian clock.

SIRT1, an anti-aging and nutrient-sensing protein, can up-regulate the expression of a number of genes involved in autophagy and energy metabolism by binding to the E-box elements in a complex with the CLOCK:BMAL1 heterodimer.^{45,46} This complex in particular creates daily oscillations in nicotinamide phosphoribosyltransferase, which is the rate-limiting enzyme in the nicotinamide adenine dinucleotide (NAD⁺) salvage pathway, thus inducing circadian oscillation of intracellular NAD⁺.^{47,48} SIRT1 can further act on circadian clock proteins by deacetylating BMAL1 and PER2, as well as other key metabolic regulators, through its regulation of NAD⁺.⁴⁸ SIRT1 histone deacetylation leads to a repressive chromatin configuration, essentially down-regulating their downstream targets.⁴⁹ As a result of SIRT1 interaction with CLOCK:BMAL1 and deacetylation of key regulator proteins, these genes have day-night rhythmicity to their transcriptional activity and temporal regulation of their target genes. Thus, the role of autophagy in the pathogenesis of

NAFLD is well described and makes these pathways a target for therapeutic interventions.^{50–52}

Circadian Clock and Glucose Homeostasis

Glucose metabolism is regulated by the circadian clock machinery. The relationship between circadian changes in metabolically important hormones (ie, glucocorticoids, insulin, ghrelin, incretins, adiponectin, and leptin) and ambient light provided the initial clues that the central clock plays an important role in glucose and nutrient homeostasis.^{53,54} Moreover, these hormones can directly affect the circadian clock in their target cells. Insulin is a regulator of *Per2* and *Rev-erba*, and both glucose and insulin can affect the expression of *Dec1* and *Dec2*.^{55,56} Low glucose/fasting induction of AMPK can repress the cryptochromes, as described in the previous section. Thus, the relationship between the circadian clock and glucose homeostasis is bidirectional.

Of particular interest, incretins, which are already the target of pharmaceuticals approved by the US Food and Drug Administration aimed to treat type 2 diabetes (T2D)




Figure 3. Circadian clock machinery. (A) In mammals, the core molecular clock comprises a series of transcription-translation feedback loops that include the transcriptional activators (CLOCK and BMAL1). The CLOCK:BMAL1 heterodimer regulates genes at a specific DNA binding sequence, the E-box, to regulate expression of hundreds of genes, including their own repressors (*Per1*, *Per2*, *Per3*, *Cry1*, and *Cry2*). Once translated, PERs and CRYs accumulate in the cytoplasm, where they are regulated by CK1 ϵ/δ and AMPK, respectively. If these PERs or CRYs are individually phosphorylated, they will undergo ubiquitylation (Ub) and proteasomal degradation. However, if PER and CRY form a heterodimer before phosphorylation by CK1 ϵ/δ , the 3-protein complex is transported to the nucleus, where it directly inhibits the CLOCK:BMAL1 heterodimer. (B) CLOCK:BMAL1 also regulates the expression of REV-ERBs and RORs. Once translated, REV-ERBs and RORs bind to the RORE DNA binding sequence, but with opposite effects. RORs promote, whereas REV-ERBs suppress, at this DNA binding sequence. Together, these clock proteins control the regulation of hundreds of genes, including the expression of *Bmal1*. (C) A third loop of the circadian clock involves DEC1 and DEC2, which are transcriptionally activated by CLOCK:BMAL1. DEC1 and DEC2 proteins migrate back into the nucleus and inhibit *Per1* transactivation by competing for E-box binding. (D) CLOCK:BMAL1 promote the transcription of *Dbp*, and REV-ERB/ROR regulate the expression of *Nfil3*. Once translated, DBP and NFIL3 promote or suppress, respectively, gene expression at the D-box DNA binding sequence. The D-box DNA binding sequence controls the expression of hundreds of clock-controlled genes.

and obesity, are being investigated as therapeutic agents in NAFLD.⁵⁷ Glucagon-like peptide 1 (GLP-1), an incretin secreted by the intestinal L cells located primarily in the ileum, is an insulin sensitizer.⁵⁸ GLP-1 release correlates with the pattern of insulin secretion, showing an increase immediately preceding the feeding period. Thus, GLP-1 contributes to the decrease in blood sugar levels by enhancing glucose-dependent insulin secretion.^{57,59} Patients with NAFLD display diminished concentrations of active incretins.^{60–62} Release of GLP-1 is circadian in rodents⁵⁹ and humans,⁶³ and this rhythmicity is lost with the consumption of a high fat/palmitate diet^{64,65} and sleep deprivation.⁶⁶ Knockdown of *Bmal1* or *Clock* results in impairment of insulin release from β -cells in response to exendin-4.^{67,68} Thus, circadian clock control of glucose-modifying hormones is not limited to the hypothalamic-pituitary axis.

Circadian clock machinery regulates glucose at a cellular level as well. Glucose homeostasis is altered in nearly all transgenic lines where the circadian clock is perturbed (Table 1).¹² The liver-specific *Bmal1*-knockout (KO) mice helped elucidate the relationship between circadian clock and GLUT2, the main hepatic glucose transporter.^{69–71} The expression of *Glut2* in mice is characterized by the peak and trough of expression corresponding to the inactive/daytime and active/nighttime states, respectively. This homeostatic stage is disrupted with loss of *Bmal1*, which results in constitutively low expression of both transcript and protein levels of *Glut2*. Hepatocyte-specific deletion of *Bmal1* in mice led to fasting hypoglycemia, hypoinsulinemia, and loss of rhythmicity in the expression of hepatic glucose metabolic genes.^{69–71}

The influence of the circadian clock is not limited to glucose transport. BMAL1 also regulates the expression of *Pgc1 α* (peroxisome proliferator-activated receptor- γ coactivator 1 α), a coactivator of gluconeogenesis.⁷² Hepatic glycogenesis is controlled by CLOCK through transcriptional activation of glycogen synthase 2 (*Gys2*), the rate-limiting enzyme in glycogen synthesis.⁷⁰ The increased expression of *Gys2* is temporally timed with the surge of postprandial glucose. *Clock*-mutant mice display severely disturbed feeding rhythms and glucose dysregulation (ie, hyperglycemia, hypoinsulinemia).^{71,73} In addition, negative mutations in *Clock* and *Bmal1* lead to impaired gluconeogenesis.⁷⁴ *Per1/Per2* double-KO mice have severe hypoglycemia in the fasting/inactive period and impaired glucose tolerance.⁶⁹ *Cry1/Cry2* double-KO mice also show glucose intolerance with elevated circulating corticosterone levels.⁷⁵ CRYs are transcriptional repressors that regulate gluconeogenic enzymes through the repression of glucocorticoid receptors⁷⁵ and CREB.^{39,76}

Not all circadian clock transgenic mice have unfavorable glucose homeostatic responses. Mice with liver-specific KO of *Ror γ* are protected against insulin resistance and hyperglycemia.⁷⁷ Incidentally, *Ror γ* -KO mice are also protected against diet-induced hepatic steatosis.³⁵ Likewise, mice with hepatic overexpression of *Cry1* had normoglycemia and improved insulin sensitivity in response to dietary challenge.³⁹ These findings have bolstered the argument for

therapeutic agents targeting circadian clock components as potential therapeutic targets for T2D and NAFLD.

Circadian Clock and Lipid and Bile Acid Homeostasis

In addition to glucose homeostasis, the circadian clock regulates lipid and bile acid metabolism. This includes regulatory transcription factors such as peroxisome proliferator-activated receptors (PPARs) and sterol regulatory element-binding proteins (SREBP) 1c (lipid metabolism) or rate-limiting enzymes such as acetyl-CoA carboxylase (ACC) (fatty acid biosynthesis) and cholesterol 7 α -hydroxylase (CYP7A1) (cholesterol and bile acid metabolism). Dysfunction in these regulatory pathways is part of the pathophysiology of NAFLD.

One of the main reasons for lipid accumulation in the hepatocytes of patients with NAFLD is dysregulation of de novo lipogenesis. SREBP1c is one of several transcription factors that can drive the induction of lipogenic genes. Overexpression of SREBP1c in the liver is associated with a 2-fold increase in alanine aminotransferase, aspartate aminotransferase, hepatic triglyceride levels, and serum free fatty acids.⁷⁸ Transgenic mice that overexpress SREBP1c displayed a fatty liver phenotype along with hyperglycemia and hyperinsulinemia.⁷⁸ The activity of this regulator is dependent on the fasting-feeding cycle and nutritional status.⁷⁹ However, more recent studies show that SREBP1c, and its targets, are also regulated by the circadian clock machinery. SREBP1c transcripts, as well as protein levels, display daily rhythmicity.⁸⁰ This rhythmicity is also reflected in the binding pattern of SREBP1c to some of its canonical target genes.⁸¹ SREBP1c is regulated by BMAL1, REV-ERB α , ROR α , ROR γ , and DEC1.^{80,82,83} For example, REV-ERB α can control hepatic cholesterol and lipid metabolism through the controlled expression of *Insig2* (insulin-induced gene 2) which is an inhibitor of SREBP1c.⁸³

Cyp7a1, the gene for the rate-limiting enzyme that converts cholesterol to bile acids, is also regulated by the circadian clock and expressed cyclically. REV-ERB α ⁸⁴ and DBP⁸⁵ regulate its transcription. *Per1/Per2* double-KO mice have abnormal expression of *Cyp7a1* and other key bile acid enzymes, suggesting that they have at least indirect regulation of bile acid synthesis.⁸⁶ Moreover, *Cyp7a1* transcription is regulated by hepatic farnesoid X receptor (FXR) and small heterodimer partner (SHP), as well as fibroblast growth factor (FGF) 15/19, which is released as a result of FXR activation in the ileum. *Fxr* and *Shp* transcription (and thus, also transcription of *Fgf15/19*) are also clock-gated.^{87,88} FXR agonists, such as obeticholic acid, are currently being investigated as potential therapeutic agents for NAFLD.⁸⁹ Finally, it should be noted that BMAL1 can further influence triglyceride metabolism through PGC1 α regulation of *Fxr*, further showing the circadian clock's multiple layers of regulatory control.⁹⁰ Altogether, the cyclic activity of CYP7A1 and its regulating enzymes generates circadian fluctuation of serum and fecal bile acids, which has implications for lipid absorption and microbiome composition.^{91,92}

Table 1. Circadian Clock Genetically Modified Lines and Their Effects on NAFLD

Mouse strain	Diet/intervention	Liver phenotype	Δ Glucose	Δ Insulin	Δ Serum triglycerides	Δ Weight	Δ Adiposity	References
Rev-erb α KO + Hepatic Rev-erb β KO	NCD	Marked hepatic steatosis with deficiency of both Rev-erbs, but only subtle changes with loss of either subtype alone	↓					177
Rev-erb α KO	NCD	Higher hepatic TG levels (trend)	↑	Normal ITT		↔	↑	178
	HFD (53% fat)	Higher hepatic TG levels (trend)	↑			↑	↑	
Per1/2 KO	NCD	Lower hepatic TG levels, depending on what time of day they were eating (oscillating)						179
Cry1/2 double KO	NCD	Hepatic steatosis	↑	↔		↓		180
	HFD (45% fat)		Impaired GTT	↑ Impaired ITT		↑	↑	
Bmal1 KO	NCD	Hepatic steatosis in global (not tissue specific) Bmal1-KO mice, but not in control mice			↑	↓		181
	HFD (60% fat)				↑	↓	↓	
Clock Δ 19 KO	NCD	Lipid engorgement of hepatocytes in Clock-KO mice compared with WT	↑	↔	↑	↑	↑	73
	HFD (45% fat)					↑	↑	
Clock Δ 19 KO	NCD	Normal				↔		182
	HFD	Lower hepatic TGs				↔		
mPer2 ^{-/-} KO	NCD and CCl ₄	Increased hepatic fibrosis in KO mice						183
Rev-erb α KO (and HDAC3 KO)	NCD	Increased hepatic TGs/steatosis in KO mice						184

NOTE. An upward arrow denotes an increase compared to control. A downward arrow denotes a decrease compared to control. A bidirectional arrow denotes no change compared to control. A lack of an arrow denotes that a comparison was not performed.

CCl₄, carbon tetrachloride; GTT, glucose tolerance test; ITT, insulin tolerance test; NCD, normal chow diet; TGs, triglycerides; WT, wild type.

Another important regulator of lipogenesis that participates in clock-controlled metabolic processes is ACC. Phosphorylation and inactivation of ACC is driven by feeding and follows a 24-hour rhythm.^{11,93} Circadian clock regulation of AMPK (eg, with ROR α) represses ACC and provides temporal control of lipogenesis.³⁴ Mice with liver-specific deletion of ACC1 exhibit lowered accumulation of hepatic triglycerides and decreased de novo fatty acid synthesis.⁹⁴ Similarly, ACC2-KO mice fed a high-fat, high-carbohydrate diet display lower body weight and less epididymal fat compared with wild-type mice.⁹⁵ Recently, an allosteric inhibitor of ACC1/2 showed promising liver-specific effects, including decreased lipogenesis, hepatic steatosis, and improved insulin sensitivity.⁹⁶

Genes involved in lipid metabolism and lipogenic processes are regulated by PPARs, a class of ligand-activated transcription factors.⁹⁷ PPAR α -KO mice fed a high-fat diet (HFD) accumulate more hepatic triglycerides and show a much higher NAFLD Activity Score.^{98,99} Conversely, activation of PPARs with fenofibrate (PPAR α) and rosiglitazone (PPAR γ) leads to amelioration of certain parameters of NAFLD, including decreased alanine aminotransferase and aspartate aminotransferase levels, steatosis, and inflammation in both mouse and clinical studies. Several randomized controlled trials have shown improvement in liver inflammation in both diabetic and nondiabetic patients with pioglitazone, another PPAR γ agonist.^{100–102} BMAL1 and CLOCK transcriptionally regulate the circadian rhythmicity of *Ppar α* . PPAR α , in turn, directly binds to PPAR response elements on the *Bmal1* promoter and promotes its expression in a positive feedback loop.^{103,104} In addition, PPAR α directly interacts with PER2 in hepatic tissue to modulate transcription of their target genes.¹⁰⁵ Because PPARs are highly circadian proteins, it is unclear whether dosing at different times of day can alter the bioavailability and effectiveness of the drug in treating NAFLD.

Circadian Clock and Endoplasmic Reticulum Stress

One of the major functions of the endoplasmic reticulum (ER) is to facilitate the correct folding of proteins. In addition, the ER is the site of lipid synthesis in hepatocytes.¹⁰⁶ Stressors such as changes in redox states or high-protein demands lead to an accumulation of unfolded or misfolded proteins, ultimately leading to ER stress. The risk of ER stress is higher at certain times of the day; thus, many of its regulatory proteins are under circadian control. As an adaptation response, the unfolded protein response (UPR), along with its component proteins, such as inositol-requiring enzyme 1 α (IRE1 α), is triggered to restore cellular homeostasis.^{107,108} The UPR plays a critical role in the pathogenesis of NAFLD.^{109–111} In mice, UPR-regulated genes exhibit a biphasic 12-hour periodicity, which is driven by circadian clock control of IRE1 α and XBP1 (X-box binding protein 1), a UPR master regulator.¹¹² In arrhythmic *Cry1/Cry2* double-KO mice, hepatic *Ire1 α* is constitutively expressed, suggesting that this pathway is under clock control. These mice also had perturbed levels of hepatic and serum triglycerides.¹¹²

Finally, the ER-associated transcription factor hepatocyte-specific CREB (CREBH) regulates stress-related lipogenesis and fatty acid oxidation in mice.^{113–115} CREBH is an important mediator of the circadian oscillator in the liver and exhibits rhythmicity in its proteolytic cleavage but not at the messenger RNA level, indicating that the circadian clock regulates it through posttranslational modification.^{115,116} CREBH is regulated indirectly by BMAL1 and AKT, and its activity is modified by DBP and NFIL3.^{116,117} Mice with CREBH deficiency have reduced levels of genes involved in fatty acid metabolism. These mice display hypertriglyceridemia and increased fatty acids, and they are more susceptible to diet-induced hepatitis and hepatic fibrosis, whereas CREBH activation protects against steatosis.^{118–120}

Luminal Content Entraines Hepatic Clock

The SCN functions as the master pacemaker and is sensitive to light signals but largely unresponsive to feeding patterns. On the other hand, the peripheral clock in the liver is dependent on feeding pattern for the amplitude and phase of the oscillation of its transcripts.^{4,121} Restricting feeding in mice to only the daytime (when they normally sleep) results in a phase shift between the central and peripheral clocks.⁸¹ Recent studies show that feeding is a stronger driver of rhythmic gene expression than circadian clock proteins, with 70% of hepatic transcripts losing rhythmicity under arrhythmic feeding.¹²² Thus, feeding is an important cue (ie, zeitgeber) for hepatic circadian clock function. It should be noted that the impetus for feeding comes from the hypothalamus, where the SCN is located. Therefore, dyssynchrony in feeding and light cycles can be considered root disorders of the SCN/hypothalamus.¹²

The gut microbiome is widely recognized for its importance in regulating metabolic physiology and as playing an important role in the pathophysiology of metabolic syndrome and NAFLD.^{123–125} Recent studies show that the gut microbiome can contribute to dysmetabolism by disruption of epithelial and hepatic circadian rhythms. Although the gut microbiome is not exposed to light, it exhibits cyclic variations that are closely intertwined to host gene expression, feeding pattern, and sex.^{126–130} Indeed, circadian disruption induced by jet lag,¹²⁶ genetic mutation of host circadian genes,^{126–128} or diet^{126,129,130} are associated with disruptions in microbe-host circadian dynamics that can result in increased adiposity, insulin resistance, and inflammation.¹² Luminal secondary (bacterially-produced) metabolites show cyclic rhythms over a 24-hour period.^{126,127,129} The cyclic variation of the bacterial taxa is one of the most robustly reproducible characteristics of the gut microbiome in murine models.¹³¹ However, the cyclic dynamics of the human gut microbiome have been difficult to study. Nevertheless, salivary samples collected over a 24-hour period^{132,133} and analysis of stool microbiome collected at different times^{134,135} suggest that the human microbiome is also highly dynamic.

Gut microbes can drive host circadian rhythms.^{12,125} Germ-free mice, antibiotic-induced microbiome-depleted

mice, or mice who lose cyclic oscillation of their gut microbiome through dietary interventions lack hepatic and intestinal circadian rhythms.^{129,136} Circadian variation of the gut microbiome can cyclically activate luminal toll-like receptors (TLRs).¹³⁶ During the rest phase, intestinal TLRs are inactive, and there is an increase in the transcription factor PPAR α , which promotes the transcription of *Rev-Erb α* . During the active phase, TLRs respond to bacterial proteins and suppress *Ppar α* transcription. With microbiome depletion, TLRs are less activated, and both PPAR α and REV-ERB α are elevated, regardless of time. Although others have reported an increase in TLR expression¹³⁷ and no changes in *Ppar α* expression,^{137,138} likely because of differences in antibiotic-depletion protocols, the observation that REV-ERB α does not cycle in microbiome-depleted mice has been reproduced in multiple studies.^{138,139} The circadian relationship between the gut microbiome and REV-ERB α was recently reproduced in another study investigating the effect of the microbiome on the circadian clock.¹³⁹ Cyclic expression of REV-ERB α leads to cyclic transcription of *Nfil3* (see Figure 3D). *Nfil3* can regulate body composition through circadian expression of the intestinal lipid metabolic program, regulating lipid absorption and its export from intestinal epithelial cells.¹³⁹ Germ-free and antibiotic-treated mice have low expression of *Nfil3* and, thus, much lower body lipid absorption.

Although these studies show a relationship between cyclic fluctuations of the gut microbiome and the circadian machinery of the host intestinal epithelial cells, it is not clear which microbially derived mediators are responsible for changes in hepatic oscillations and, thus, metabolic functions. Two potential mediators are bile acids and short-chain fatty acids (SCFAs) (Supplementary Figure 1).¹²⁵ As discussed earlier, bile acids and key enzymes in their production and uptake are cyclically expressed and regulated by circadian clock genes.^{84,91,140} Although incompletely characterized, secondary bile acids can affect intestinal and hepatic circadian gene expression in vivo.¹⁴¹ Increased expression of bile salt hydrolase, a bacterial enzyme that deconjugates bile acids and allows for the production of secondary bile acids, induces a shift in ileal and hepatic circadian gene expression in microbiome-depleted mice.¹⁴² Moreover, sleep disruption, altered feeding pattern, and circadian gene KOs alter the expression of key bile acid regulatory genes.^{85,91} Cecal SCFAs have cyclic fluctuations that are perturbed with a HFD.¹²⁹ Although the mechanism is incompletely understood, supplementation with SCFAs can cause changes in *Bmal1* and *Per2* gene expression. Interestingly, *Bmal1*-KO mice lose cyclic fluctuation of their fecal SCFAs.¹⁴³ Overall, these studies suggest a reciprocal interaction between the circadian clock and luminal bile acids and SCFAs.

Dyssynchrony and Metabolic Syndrome

Synchrony between the SCN and hepatic circadian clock temporally organizes the expression of a large number of metabolic regulatory genes to the daily pattern of food availability. In the absence of a robust hepatic circadian clock, the organism becomes susceptible to various

metabolic disorders, including increased adiposity, ectopic steatosis, and insulin resistance.¹² Genetically modified mouse lines affecting various clock genes show the interconnectedness between circadian rhythms and metabolic regulators. Circadian gene transgenic lines nearly universally have perturbed metabolic phenotypes (eg, insulin resistance, increased adiposity) (Table 1).¹² Moreover, nearly all dietary-induced dysmetabolic mouse models have perturbed feeding patterns, often characterized by the loss of discrete meals and the spread of caloric intake throughout the day and night.¹² Thus, circadian dyssynchrony, characterized by phase shifts and dampening of circadian gene oscillations that lead to misalignment of peripheral and central circadian rhythms, is common to nearly all models of metabolic syndrome, including NAFLD.

Murine Models

Efforts to decipher the mechanistic relationship between circadian clock and host metabolism have been quite fruitful, especially over the last two decades. In general, decreased expression of the clock genes in the positive limb result in worse metabolic outcomes. Conversely, decreased expression of the negative limb (see Figure 2) clock genes leads to improved metabolic outcomes and protection from hepatic steatosis. A summary of selected studies of clock transgenic mice and their metabolic characteristics are presented in Table 1.

Environmental stressors, such as changes in diet, sleeping pattern, or lighting conditions, can also cause circadian dysynchrony. For example, in the diet-induced obesity (DIO) model—where mice are given an HFD and develop increased adiposity, hyperlipidemia, steatosis, and insulin resistance—mice spread their caloric intake throughout the day and night, thereby doubling the percentage of food eaten during the day/inactive period.¹⁴⁴ This altered feeding pattern causes a disruption of the hepatic circadian clock. Leptin-KO (*Ob/Ob*) mice, a genetic model of obesity characterized by hyperphagia due to leptin deficiency, also have dampened oscillation of their circadian clock before weight gain.^{145,146} In addition to feeding, changes to light (eg, light at night) or disruption of sleep in mice also leads to dysmetabolic states characterized by increased features of metabolic syndrome, including steatosis.¹⁶

Human Studies

As in mice, human studies show a strong relationship between the circadian dyssynchrony and dysmetabolism. Genetic studies have been instrumental in showing this relationship. Haplotype analysis of patients with NAFLD showed an association between steatosis and CLOCK, where gene variants have a potential role in the susceptibility, pathogenesis, and disease progression.¹⁴⁷ A similar study reported the putative role of CLOCK polymorphisms, which were associated with a 1.8-fold increase in susceptibility to obesity.¹⁴⁸ SNPs in CLOCK have also been associated with metabolic syndrome, hypoglycemia, obesity, and T2D.¹⁴⁹ Genome-wide association studies provide strong evidence for the role of circadian gene variants in susceptibility to

Table 2. Selected Studies Investigating Associations Between Sleep or Food Behaviors and Metabolic Syndrome or NAFLD

Article	Study type	Population	Control	Comparator	Key findings
Selected studies investigating associations between sleep disturbance or TRF and metabolic syndrome					
Spiegel et al (1999) ¹⁸⁵	Observation study	Healthy volunteers (N = 11)	N/A	Sleep deprivation	• Sleep-deprivation reduced glucose tolerance and evening cortisol concentrations.
Wang et al (2014) ¹⁸⁶	Meta-analysis of 13 observational studies	Workers exposed to night shift (N = 2286)	Day workers	Shift workers	• Higher risk of metabolic syndrome in workers exposed to night shift (OR, 1.5–1.6).
Vyas et al (2012) ¹⁸⁷	Meta-analysis of 9 case-control studies, 11 prospective cohort studies, and 6 retrospective cohort studies	Workers exposed to shift work (N = 2,000,000)	Day workers	Shift workers	• Shift work is associated with an increased risk of myocardial infarction and coronary events (OR, 1.23).
Karlsson et al (2001) ¹⁸⁸	Observational study	Working adults (N = 27,485)	Day workers	Shift workers	• OR of obesity was approximately 1.3 higher in shift workers compared with day workers. • Dyslipidemia clustered with shift workers compared with day workers ($P < .05$).
Wilkinson et al (2019) ¹⁸⁹	Paired-sample trial	Patients with metabolic syndrome (N = 19)	2-week baseline of eating interval of ≥ 14 h per day	12-weeks of time-restricted eating (TRE) to a 10-hour interval	• After 12 weeks of TRF, there was a 3% reduction in weight ($P = .0003$), 11% reduction in LDL-C ($P = .016$), and 8% reduction in estimated calorie intake ($P = .007$) but no difference in insulin resistance ($P = .107$).
Selected studies investigating associations between sleep disturbances and/or shift work and NAFLD					
Hsieh et al (2011) ¹⁹⁰	Observational	All-male Tokyo office workers (N = 8157)	Sleep >7 hours	Sleep <7 hours	• Rates of NAFLD were higher in patients who slept <7 hours (OR, 1.23).
Kim et al (2013) ¹⁹¹	Observational	Workers and spouses at a Korean conglomerate (N = 69,463)	Sleep >5 hours	Sleep <5 hours	• Rates of NAFLD were higher among women (but not men) who slept <5 hours (OR, 1.59 in women; 1.03 in men).
Imaizumi et al (2015) ¹⁹²	Observational studies	General patients (N = 3986)	Sleep duration 6 to ≤ 7 h	Sleep duration ≤ 6 hours	• Rates of NAFLD were higher among women (but not men) who slept ≤ 6 hours (OR, 1.44 in women; 0.98 in men).
Bernsmeier et al (2015) ¹⁹³	Case-control study	Cohort in Basel (N = 68)	Healthy volunteers (n = 22)	Biopsy-proven NAFLD group (n = 46)	• NAFLD group had shorter sleep duration and poorer sleep quality compared with control group ($P < .01$). In patients with NAFLD, but not in control individuals, daytime sleepiness was correlated with higher liver enzyme levels and insulin resistance. • Patients with NAFLD with fibrosis had stronger correlation to daytime sleepiness ($P = .02$).
Marin-Alejandro et al (2019) ¹⁹⁴	Cross-sectional study	Fatty Liver in Obesity participants in Spain (n = 134)	Normal-weight individuals without NAFLD (n = 40)	Overweight patients with NAFLD (n 9 4)	• Adjusted odds of sleep disturbance was higher in NAFLD group (OR, 1.59). • Regression models predicted approximately 20% of variability in liver stiffness could be attributed to sleep disturbance or sleep quality.
Liu et al (2016) ¹⁹⁵	Cohort study	Patients who developed NAFLD after 5 years of follow-up (n = 2197)	Sleep duration 8–9 hours, sleep duration >9 hours	Sleep duration of 7–8 hours	• Adjusted odds of NAFLD was higher with longer sleep durations (OR, 1.21 for the 8–9-hours group; 1.31 in >9 hours group).

Table 2. Continued

Article	Study type	Population	Control	Comparator	Key findings
Balakrishnan et al (2017) ¹⁹⁶	Observational study	General workers (n = 8159)	Non-shift workers	Shift workers	• Adjusted odds of NAFLD did not differ between shift workers and non-shift workers (OR, 1.11).
Katsagoni et al (2018) ¹⁹⁷	Randomized controlled trial	Patients with NAFLD (n = 63)	Control individuals without intervention	Mediterranean lifestyle (adequate sleep and activity)	• Mediterranean lifestyle was associated with a decrease in liver stiffness (adjusted OR, 0.78).
Katsagoni et al (2017) ¹⁹⁸	Case-control study	Patients with NAFLD and control individuals (n = 155)	Control individuals without NAFLD (n = 55)	Patients with NAFLD (n = 100)	• Optimal sleep duration was inversely associated with NAFLD presence (OR, 0.38; <i>P</i> = .05).
Selected studies investigating link between OSA and NAFLD					
Musso et al (2013) ¹⁹⁹	Meta-analysis of 18 cross-sectional studies	18 cross-sectional studies (n = 2183) of patients with OSA who were assessed for NAFLD	Patients with OSA without NAFLD	Patients with OSA with NAFLD	• OSA was associated with nonalcoholic steatohepatitis (OR, 2.37) and advanced fibrosis (OR, 2.3) in patients with biopsy-proven NAFLD.
Pamidi and Tasali (2012) ²⁰⁰	Observational	Healthy and young patients with OSA	Control individuals without OSA (n = 20)	Patients with OSA (n = 12)	• OSA was associated with a 27% lower insulin sensitivity compared with control individuals.

LDL-C, low-density lipoprotein cholesterol; N/A, not applicable; OR, odds ratio; OSA, obstructive sleep apnea.

metabolic syndrome. They have shown links between PER3 and BMAL1 and T2D, BMAL1 and hypertension, and CRY2 and insulin resistance.¹⁵⁰

Environmental and behavioral effects on the circadian clock also have metabolic consequences, which are summarized in Table 2. Although human studies show significant association between sleep disturbances, shift work, and genetic predispositions to metabolic syndrome, it is not clear whether the mechanisms driving these metabolic phenotypes are similar to what has been observed in mice. Provocatively, one of the studies that investigated the circadian dynamics of the gut microbiome suggests that the microbiota could be mediating the dysmetabolic phenotype of circadian dyssynchrony in humans.¹²⁶ Gnotobiotic mice that received a fecal transplant from jet-lagged individuals had increased adiposity and insulin resistance compared with gnotobiotic mice who received pre-jet lag and post-jet lag stool from the same individuals.¹²⁶ However, only 2 patients were used in this small cohort, and more vigorous investigation of the role of the gut microbiome in circadian dyssynchrony is necessary.

Correction of Dyssynchrony With Time-Restricted Feeding

It was not clear until recently whether correcting circadian dyssynchrony is sufficient to reverse the dysmetabolic effects of these various insults. TRF, a behavioral paradigm where feeding is consolidated to the active period, aligns peripheral and central circadian rhythms.¹⁵¹ Correction of circadian dyssynchrony with TRF prevents and treats the metabolic consequences of a large variety of insults.^{12,152–155} It should be noted that TRF is not synonymous with intermittent fasting; the former is focused on aligning peripheral and central circadian rhythms. When applied to human populations, the number of calories consumed is irrelevant in the time-restricted diet, which is part of its appeal. Intermittent fasting, on the other hand, focuses on creating periods of fasting, usually on alternate days, to induce changes to metabolism. Caloric intake is strictly regulated on fasting days.

Mice in the TRF condition have access to food only in the nocturnal (active) phase, which promotes natural feeding rhythms and restores the amplitude and phase of oscillations of peripheral circadian clock genes, which are obliterated in DIO mice.¹⁵¹ TRF also restores cycling of metabolic regulators (eg, AMPK, mTOR, and CREB) and their downstream targets. The metabolic effects of restoring synchrony can be profound. Although TRF mice consume the same number of calories and have similar activity levels, they do not have the dysmetabolic phenotype of DIO mice. TRF prevents and treats the increased adiposity, insulin and leptin resistance, steatohepatitis, and dyslipidemia that occur with the DIO condition.^{130,151,156,157} Subsequent studies show that TRF protects against the metabolic effects of a variety of dysmetabolic diets, including the Western diet, high-fructose, and high-sucrose diets.¹⁵⁶

TRF also protects against the dysmetabolism induced by circadian gene KOs. *Per1*-, hepatic *Bmal1*-, hepatic *Rev-Erba*/

β -, and *Cry1/Cry2*-KO mice all have dysmetabolic phenotypes when given free access to HFD. In all cases, TRF prevented the dysmetabolic phenotype.⁸² TRF reduces hepatic steatosis and enhances the cellular response to metabolic stress. Hence, the benefits from TRF likely arise from restoring daily feeding/fasting rhythms and balancing nutrient absorptions with cellular stress response. TRF also protected against the dysmetabolic phenotypes due to aging, obesity, and central circadian rhythm dysfunction in *Drosophila*.^{158,159}

Contrary to popular belief, instead of consuming three distinct meals, humans consume calories sporadically and over a large range of time within a 24-hour period.¹⁶⁰ Thus, TRF could potentially be an easily adoptable behavioral intervention that improves outcomes in patients with metabolic syndrome who have irregular eating habits. Several small-scale human TRF studies have shown interesting although sometimes contradictory results. In one randomized crossover trial, healthy volunteers were first assigned to either a standard diet of three meals per day or to receive one meal in the evening, both without caloric restriction.¹⁶¹ The two groups did not have a difference in the number of calories consumed. The group with one meal a day had higher morning fasting plasma glucose levels and a delayed insulin response in an oral glucose tolerance test. A different study compared healthy male participants consuming a standard diet of three meals per day with a group with TRF in the afternoon to evening. In this case, the TRF group had decreased fat mass, without any difference in triglycerides, total cholesterol, or low-density lipoprotein.¹⁶² A subsequent randomized, supervised controlled feeding trial in humans showed that participants in the early TRF group had improved insulin sensitivity, appetite, and blood pressure but had higher levels of triglycerides and total cholesterol.¹⁶³ A more recent study aimed to compare early vs late TRF among men at risk for T2D (monitored with continuous glucose monitors) found that the early-TRF group had lower mean fasting glucose, whereas both TRF groups had an improvement in glucose response to meals.¹⁶⁴ However, it is not clear whether the investigators controlled for the length of time the participants were fasting during their sample collection, necessitating additional larger studies. A recent study investigated the effects of TRF on patients with metabolic syndrome undergoing pharmacotherapy. In this single-arm, paired-sample trial of 19 patients who had an initial eating window of >14 hours and were undergoing statin or anti-hypertensive therapy, imposing a 10-hour feeding window (monitored with continuous glucose monitors) led to improved cardiometabolic health. Although these studies are promising, a study investigating the effects of TRF on NAFLD endpoints has not yet been conducted.

Conclusion

NAFLD is a complex disease that is associated with a multitude of metabolic perturbations.^{7,165} Because many circadian clock-controlled genes are vital participants in metabolic processes of the body, it is not surprising that some of these rhythmic genes can be potential targets for therapy. Behavioral interventions, such as TRF, may have

benefits in NAFLD that are independent of its weight loss effects. TRF may be easier for patients to adopt because it does not restrict calories or require a specific macronutrient dietary profile.¹⁶⁰

Chemical modulators of clock-regulated processes such as bile acid metabolism (FXR agonist obeticholic acid, FGF15/19 analogue NGM282), incretin response (GLP-1 receptor agonist liraglutide), and PPAR-controlled lipogenic mechanisms (PPAR α activator elafibranor) are under various stages of the clinical trial process.^{11,166,167} However, because some components of the core clock directly participate in metabolic processes and the oscillator itself is responsive to resetting stimuli, small molecules directed toward clock components might provide alternative targets for therapeutic intervention.^{47,168} In this light, administration of compound KL001, an activator of CRY proteins, improved glucose tolerance in DIO mice.^{169,170} This is consistent with the known role of CRYs in the inhibition of gluconeogenesis.⁷⁵ Recent studies show that CRY1 is targeted for degradation by autophagy, thus depressing hepatic gluconeogenesis in a temporal manner.⁷⁶ Further research into these sites on CRY1 might direct researchers toward novel drug targets for the treatment of hyperglycemia and insulin resistance. Nobiletin, a drug that enhances the amplitude of *Ror* expression, prevents weight gain and improves metabolic parameters in DIO and *Ob/Ob* mice.^{171,172} Additionally, treatment with REV-ERB agonists SR9009 and SR9011 led to improved metabolic endpoints in DIO mice.^{173,174} These drugs could potentially be beneficial in NAFLD.

Furthermore, the circadian clock extends its functional network to the control of global physiological processes, including the targets of nearly all drugs affecting metabolic syndrome spectrum disorders.¹⁷⁵ The rhythmicity imparted by the circadian clock in drug metabolic and detoxification processes can be a potential target for the treatment of NAFLD.^{10,11} By better accounting for the circadian cycling of their targets, new drugs can be administered in a way that maximizes benefits, and minimizes adverse effects, potentially arriving at a therapeutic dose with fewer administrations and lower dosages.¹⁷⁶ Further study of the chronopharmacology and circadian xenobiotics of small-molecule therapeutics might provide better insight into the proper temporal profile and contribute to improved efficacy.

Supplementary Material

Note: To access the supplementary material accompanying this article, visit the online version of *Gastroenterology* at www.gastrojournal.org, and at <https://doi.org/10.1053/j.gastro.2020.01.050>.

References

1. Gachon F, Nagoshi E, Brown SA, et al. The mammalian circadian timing system: from gene expression to physiology. *Chromosoma* 2004;113:103–112.
2. Yamazaki S, Numano R, Abe M, et al. Resetting central and peripheral circadian oscillators in transgenic rats. *Science* 2000;288(5466):682–685.

3. Mure LS, Le HD, Benegiamo G, et al. Diurnal transcriptome atlas of a primate across major neural and peripheral tissues. *Science* 2018;359(6381):eaao0318.
4. Damiola F, Le Minh N, Preitner N, et al. Restricted feeding uncouples circadian oscillators in peripheral tissues from the central pacemaker in the suprachiasmatic nucleus. *Genes Dev* 2000;14:2950–2961.
5. Buijs R, Salgado R, Sabath E, et al. Peripheral circadian oscillators: time and food. *Prog Mol Biol Transl Sci* 2013;119:83–103.
6. Shetty A, Hsu JW, Manka PP, et al. Role of the circadian clock in the metabolic syndrome and nonalcoholic fatty liver disease. *Dig Dis Sci* 2018;63:3187–3206.
7. Shi D, Chen J, Wang J, et al. Circadian clock genes in the metabolism of non-alcoholic fatty liver disease. *Front Physiol* 2019;10:423.
8. Gnocchi D, Custodero C, Sabba C, et al. Circadian rhythms: a possible new player in non-alcoholic fatty liver disease pathophysiology. *J Mol Med (Berl)* 2019;97:741–759.
9. Reinke H, Asher G. Circadian clock control of liver metabolic functions. *Gastroenterology* 2016;150:574–580.
10. Tahara Y, Shibata S. Circadian rhythms of liver physiology and disease: experimental and clinical evidence. *Nat Rev Gastroenterol Hepatol* 2016;13:217–226.
11. Mukherji A, Bailey SM, Staels B, et al. The circadian clock and liver function in health and disease. *J Hepatol* 2019;71:200–211.
12. Zarrinpar A, Chaix A, Panda S. Daily eating patterns and their impact on health and disease. *Trends Endocrinol Metab* 2016;27:69–83.
13. Cederroth CR, Albrecht U, Bass J, et al. Medicine in the fourth dimension. *Cell Metab* 2019;30:238–250.
14. Challet E. The circadian regulation of food intake. *Nat Rev Endocrinol* 2019;15:393–405.
15. Panda S. Circadian physiology of metabolism. *Science* 2016;354(6315):1008–1015.
16. Stenvers DJ, Scheer F, Schrauwen P, et al. Circadian clocks and insulin resistance. *Nat Rev Endocrinol* 2019;15:75–89.
17. Wang Z, Wu Y, Li L, et al. Intermolecular recognition revealed by the complex structure of human CLOCK-BMAL1 basic helix-loop-helix domains with E-box DNA. *Cell Res* 2013;23:213–224.
18. Yan J, Wang H, Liu Y, et al. Analysis of gene regulatory networks in the mammalian circadian rhythm. *PLoS Comput Biol* 2008;4(10):e1000193.
19. Knippschild U, Gocht A, Wolff S, et al. The casein kinase 1 family: participation in multiple cellular processes in eukaryotes. *Cell Signal* 2005;17:675–689.
20. Busino L, Bassemann F, Maiolica A, et al. SCF^{Fbxl3} controls the oscillation of the circadian clock by directing the degradation of cryptochrome proteins. *Science* 2007;316(5826):900–904.
21. Vielhaber E, Eide E, Rivers A, et al. Nuclear entry of the circadian regulator mPER1 is controlled by mammalian casein kinase I epsilon. *Mol Cell Biol* 2000;20:4888–4899.
22. Zhang Y, Fang B, Emmett MJ, et al. Discrete functions of nuclear receptor Rev-erb α couple metabolism to the clock. *Science* 2015;348(6242):1488–1492.
23. Sato TK, Panda S, Miraglia LJ, et al. A functional genomics strategy reveals Rora as a component of the mammalian circadian clock. *Neuron* 2004;43:527–537.
24. Preitner N, Damiola F, Lopez-Molina L, et al. The orphan nuclear receptor REV-ERB α controls circadian transcription within the positive limb of the mammalian circadian oscillator. *Cell* 2002;110:251–260.
25. Everett LJ, Lazar MA. Nuclear receptor Rev-erb α : up, down, and all around. *Trends Endocrinol Metab* 2014;25:586–592.
26. Akashi M, Takumi T. The orphan nuclear receptor ROR α regulates circadian transcription of the mammalian core-clock *Bmal1*. *Nat Struct Mol Biol* 2005;12:441–448.
27. Guillaumond F, Dardente H, Giguère V, et al. Differential control of Bmal1 circadian transcription by REV-ERB and ROR nuclear receptors. *J Biol Rhythms* 2005;20:391–403.
28. Kato Y, Kawamoto T, Fujimoto K, Noshiro M. DEC1/STRA13/SHARP2 and DEC2/SHARP1 coordinate physiological processes, including circadian rhythms in response to environmental stimuli. *Curr Top Dev Biol* 2014;110:339–372.
29. Honma S, Kawamoto T, Takagi Y, et al. *Dec1* and *Dec2* are regulators of the mammalian molecular clock. *Nature* 2002;419(6909):841–844.
30. Yamajuku D, Shibata Y, Kitazawa M, et al. Cellular DBP and E4BP4 proteins are critical for determining the period length of the circadian oscillator. *FEBS Lett* 2011;585:2217–2222.
31. Yoshitane H, Asano Y, Sagami A, et al. Functional D-box sequences reset the circadian clock and drive mRNA rhythms. *Commun Biol* 2019;2:300.
32. Duffield GE. DNA microarray analyses of circadian timing: the genomic basis of biological time. *J Neuroendocrinol* 2003;15:991–1002.
33. Zhang R, Lahens NF, Ballance HI, et al. A circadian gene expression atlas in mammals: implications for biology and medicine. *Proc Natl Acad Sci U S A* 2014;111:16219–16224.
34. Lamia KA, Sachdeva UM, DiTacchio L, et al. AMPK regulates the circadian clock by cryptochrome phosphorylation and degradation. *Science* 2009;326(5951):437–440.
35. Kim EJ, Yoon YS, Hong S, et al. Retinoic acid receptor-related orphan receptor α -induced activation of adenosine monophosphate-activated protein kinase results in attenuation of hepatic steatosis. *Hepatology* 2012;55:1379–1388.
36. Travnickova-Bendova Z, Cermakian N, Reppert SM, et al. Bimodal regulation of *mPeriod* promoters by CREB-dependent signaling and CLOCK/BMAL1 activity. *Proc Natl Acad Sci U S A* 2002;99:7728–7733.
37. O'Neill JS, Maywood ES, Chesham JE, et al. cAMP-dependent signaling as a core component of the mammalian circadian pacemaker. *Science* 2008;320(5878):949–953.

38. Sun X, Dang F, Zhang D, et al. Glucagon-CREB/CRTC2 signaling cascade regulates hepatic BMAL1 protein. *J Biol Chem* 2015;290:2189–2197.
39. Zhang EE, Liu Y, Dentin R, et al. Cryptochrome mediates circadian regulation of cAMP signaling and hepatic gluconeogenesis. *Nat Med* 2010;16:1152–1156.
40. Kalsbeek A, la Fleur S, Fliers E. Circadian control of glucose metabolism. *Mol Metab* 2014;3:372–383.
41. Luciano AK, Zhou W, Santana JM, et al. CLOCK phosphorylation by AKT regulates its nuclear accumulation and circadian gene expression in peripheral tissues. *J Biol Chem* 2018;293:9126–9136.
42. Rabinowitz JD, White E. Autophagy and metabolism. *Science* 2010;330(6009):1344–1348.
43. Kim J, Kundu M, Viollet B, et al. AMPK and mTOR regulate autophagy through direct phosphorylation of Ulk1. *Nat Cell Biol* 2011;13:132–141.
44. Khapre RV, Kondratova AA, Patel S, et al. BMAL1-dependent regulation of the mTOR signaling pathway delays aging. *Aging (Albany NY)* 2014;6:48–57.
45. Nakahata Y, Sahar S, Astarita G, et al. Circadian control of the NAD⁺ salvage pathway by CLOCK-SIRT1. *Science* 2009;324(5927):654–657.
46. Eckel-Mahan K, Sassone-Corsi P. Metabolism and the circadian clock converge. *Physiol Rev* 2013;93:107–135.
47. Asher G, Schibler U. Crosstalk between components of circadian and metabolic cycles in mammals. *Cell Metab* 2011;13:125–137.
48. Zhou B, Zhang Y, Zhang F, et al. CLOCK/BMAL1 regulates circadian change of mouse hepatic insulin sensitivity by SIRT1. *Hepatology* 2014;59:2196–2206.
49. Nakahata Y, Kaluzova M, Grimaldi B, et al. The NAD⁺-dependent deacetylase SIRT1 modulates CLOCK-mediated chromatin remodeling and circadian control. *Cell* 2008;134:329–340.
50. Tanaka S, Hikita H, Tatsumi T, et al. Rubicon inhibits autophagy and accelerates hepatocyte apoptosis and lipid accumulation in nonalcoholic fatty liver disease in mice. *Hepatology* 2016;64:1994–2014.
51. Lin CW, Zhang H, Li M, et al. Pharmacological promotion of autophagy alleviates steatosis and injury in alcoholic and non-alcoholic fatty liver conditions in mice. *J Hepatol* 2013;58:993–999.
52. Fucho R, Martinez L, Baulies A, et al. ASMase regulates autophagy and lysosomal membrane permeabilization and its inhibition prevents early stage non-alcoholic steatohepatitis. *J Hepatol* 2014;61:1126–1134.
53. Albrecht U. Timing to perfection: the biology of central and peripheral circadian clocks. *Neuron* 2012;74:246–260.
54. Van Cauter E. Sleep disturbances and insulin resistance. *Diabet Med* 2011;28:1455–1462.
55. Tahara Y, Otsuka M, Fuse Y, et al. Refeeding after fasting elicits insulin-dependent regulation of *Per2* and *Rev-erb α* with shifts in the liver clock. *J Biol Rhythms* 2011;26:230–240.
56. Sato F, Muragaki Y, Kawamoto T, et al. Rhythmic expression of DEC2 protein in vitro and in vivo. *Biomed Rep* 2016;4:704–710.
57. Zarrinpar A, Loomba R. Review article: the emerging interplay among the gastrointestinal tract, bile acids and incretins in the pathogenesis of diabetes and non-alcoholic fatty liver disease. *Aliment Pharmacol Ther* 2012;36:909–921.
58. Holst JJ. The physiology of glucagon-like peptide 1. *Physiol Rev* 2007;87:1409–1439.
59. Gil-Lozano M, Mingomataj EL, Wu WK, et al. Circadian secretion of the intestinal hormone GLP-1 by the rodent L cell. *Diabetes* 2014;63:3674–3685.
60. Seghieri M, Christensen AS, Andersen A, et al. Future perspectives on GLP-1 receptor agonists and GLP-1/glucagon receptor co-agonists in the treatment of NAFLD. *Front Endocrinol (Lausanne)* 2018;9:649.
61. Bernsmeier C, Meyer-Gerspach AC, Blaser LS, et al. Glucose-induced glucagon-like peptide 1 secretion is deficient in patients with non-alcoholic fatty liver disease. *PLoS One* 2014;9(1):e87488.
62. Firmeisz G, Varga T, Lengyel G, et al. Serum dipeptidyl peptidase-4 activity in insulin resistant patients with non-alcoholic fatty liver disease: a novel liver disease biomarker. *PLoS One* 2010;5(8):e12226.
63. Galindo Munoz JS, Jimenez Rodriguez D, Hernandez Morante JJ. Diurnal rhythms of plasma GLP-1 levels in normal and overweight/obese subjects: lack of effect of weight loss. *J Physiol Biochem* 2015;71:17–28.
64. Martchenko A, Oh RH, Wheeler SE, et al. Suppression of circadian secretion of glucagon-like peptide-1 by the saturated fatty acid, palmitate. *Acta Physiol (Oxf)* 2018;222(4):e13007.
65. Gil-Lozano M, Wu WK, Martchenko A, et al. High-fat diet and palmitate alter the rhythmic secretion of glucagon-like peptide-1 by the rodent L-cell. *Endocrinology* 2016;157:586–599.
66. Gil-Lozano M, Hunter PM, Behan LA, et al. Short-term sleep deprivation with nocturnal light exposure alters time-dependent glucagon-like peptide-1 and insulin secretion in male volunteers. *Am J Physiol Endocrinol Metab* 2016;310(1):E41–E50.
67. Marcheva B, Ramsey KM, Buhr ED, et al. Disruption of the clock components CLOCK and BMAL1 leads to hypoinsulinaemia and diabetes. *Nature* 2010;466(7306):627–631.
68. Brubaker PL, Gil-Lozano M. Glucagon-like peptide-1: the missing link in the metabolic clock? *J Diabetes Investig* 2016;7(Suppl 1):70–75.
69. Lamia KA, Storch KF, Weitz CJ. Physiological significance of a peripheral tissue circadian clock. *Proc Natl Acad Sci U S A* 2008;105:15172–15177.
70. Frese T, Bazwinsky I, Muhlbauer E, et al. Circadian and age-dependent expression patterns of GLUT2 and glucokinase in the pancreatic β -cell of diabetic and nondiabetic rats. *Horm Metab Res* 2007;39:567–574.
71. Doi R, Oishi K, Ishida N. CLOCK regulates circadian rhythms of hepatic glycogen synthesis through transcriptional activation of *Gys2*. *J Biol Chem* 2010;285:22114–22121.
72. Liu C, Li S, Liu T, et al. Transcriptional coactivator PGC-1 α integrates the mammalian clock and energy metabolism. *Nature* 2007;447(7143):477–481.

73. Turek FW, Joshu C, Kohsaka A, et al. Obesity and metabolic syndrome in circadian *Clock* mutant mice. *Science* 2005;308(5724):1043–1045.
74. Rudic RD, McNamara P, Curtis AM, et al. BMAL1 and CLOCK, two essential components of the circadian clock, are involved in glucose homeostasis. *PLoS Biol* 2004;2(11):e377.
75. Lamia KA, Papp SJ, Yu RT, et al. Cryptochromes mediate rhythmic repression of the glucocorticoid receptor. *Nature* 2011;480(7378):552–556.
76. Toledo M, Batista-Gonzalez A, Merheb E, et al. Autophagy regulates the liver clock and glucose metabolism by degrading CRY1. *Cell Metab* 2018;28:268–281.
77. Takeda Y, Kang HS, Freudenberg J, et al. Retinoic acid-related orphan receptor γ (ROR γ): a novel participant in the diurnal regulation of hepatic gluconeogenesis and insulin sensitivity. *PLoS Genet* 2014;10(5):e1004331.
78. Knebel B, Haas J, Hartwig S, et al. Liver-specific expression of transcriptionally active SREBP-1c is associated with fatty liver and increased visceral fat mass. *PLoS One* 2012;7(2):e31812.
79. Li Y, Xu S, Mihaylova MM, et al. AMPK phosphorylates and inhibits SREBP activity to attenuate hepatic steatosis and atherosclerosis in diet-induced insulin-resistant mice. *Cell Metab* 2011;13:376–388.
80. Zhang Y, Papazyan R, Damle M, et al. The hepatic circadian clock fine-tunes the lipogenic response to feeding through ROR α/γ . *Genes Dev* 2017;31:1202–1211.
81. Vollmers C, Gill S, DiTacchio L, et al. Time of feeding and the intrinsic circadian clock drive rhythms in hepatic gene expression. *Proc Natl Acad Sci U S A* 2009;106:21453–21458.
82. Chaix A, Lin T, Le HD, et al. Time-restricted feeding prevents obesity and metabolic syndrome in mice lacking a circadian clock. *Cell Metab* 2019;29:303–319.
83. Le Martelot G, Claudel T, Gatfield D, et al. REV-ERB α participates in circadian SREBP signaling and bile acid homeostasis. *PLoS Biol* 2009;7(9):e1000181.
84. Duez H, van der Veen JN, Duhem C, et al. Regulation of bile acid synthesis by the nuclear receptor Rev-erb α . *Gastroenterology* 2008;135:689–698.
85. Ferrell JM, Chiang JY. Short-term circadian disruption impairs bile acid and lipid homeostasis in mice. *Cell Mol Gastroenterol Hepatol* 2015;1:664–677.
86. Ma K, Xiao R, Tseng HT, et al. Circadian dysregulation disrupts bile acid homeostasis. *PLoS One* 2009;4(8):e6843.
87. Yang X, Downes M, Yu RT, et al. Nuclear receptor expression links the circadian clock to metabolism. *Cell* 2006;126:801–810.
88. Stroeve JH, Brufau G, Stellaard F, et al. Intestinal FXR-mediated FGF15 production contributes to diurnal control of hepatic bile acid synthesis in mice. *Lab Invest* 2010;90:1457–1467.
89. Neuschwander-Tetri BA, Loomba R, Sanyal AJ, et al. Farnesoid X nuclear receptor ligand obeticholic acid for non-cirrhotic, non-alcoholic steatohepatitis (FLINT): a multicentre, randomised, placebo-controlled trial. *Lancet* 2015;385(9972):956–965.
90. Zhang Y, Castellani LW, Sinal CJ, et al. Peroxisome proliferator-activated receptor-gamma coactivator 1 α (PGC-1 α) regulates triglyceride metabolism by activation of the nuclear receptor FXR. *Genes Dev* 2004;18:157–169.
91. Zhang YK, Guo GL, Klaassen CD. Diurnal variations of mouse plasma and hepatic bile acid concentrations as well as expression of biosynthetic enzymes and transporters. *PLoS One* 2011;6(2):e16683.
92. Galman C, Angelin B, Rudling M. Bile acid synthesis in humans has a rapid diurnal variation that is asynchronous with cholesterol synthesis. *Gastroenterology* 2005;129:1445–1453.
93. Davies SP, Carling D, Munday MR, et al. Diurnal rhythm of phosphorylation of rat liver acetyl-CoA carboxylase by the AMP-activated protein kinase, demonstrated using freeze-clamping. Effects of high fat diets. *Eur J Biochem* 1992;203:615–623.
94. Mao J, DeMayo FJ, Li H, et al. Liver-specific deletion of acetyl-CoA carboxylase 1 reduces hepatic triglyceride accumulation without affecting glucose homeostasis. *Proc Natl Acad Sci U S A* 2006;103:8552–8557.
95. Abu-Elheiga L, Wu H, Gu Z, et al. Acetyl-CoA carboxylase 2^{-/-} mutant mice are protected against fatty liver under high-fat, high-carbohydrate dietary and de novo lipogenic conditions. *J Biol Chem* 2012;287:12578–12588.
96. Goedeke L, Bates J, Vatner DF, et al. Acetyl-CoA carboxylase inhibition reverses NAFLD and hepatic insulin resistance but promotes hypertriglyceridemia in rodents. *Hepatology* 2018;68:2197–2211.
97. Pawlak M, Lefebvre P, Staels B. Molecular mechanism of PPAR α action and its impact on lipid metabolism, inflammation and fibrosis in non-alcoholic fatty liver disease. *J Hepatol* 2015;62:720–733.
98. Stienstra R, Mandart S, Patsouris D, et al. Peroxisome proliferator-activated receptor α protects against obesity-induced hepatic inflammation. *Endocrinology* 2007;148:2753–2763.
99. Patsouris D, Reddy JK, Muller M, et al. Peroxisome proliferator-activated receptor α mediates the effects of high-fat diet on hepatic gene expression. *Endocrinology* 2006;147:1508–1516.
100. Aithal GP, Thomas JA, Kaye PV, et al. Randomized, placebo-controlled trial of pioglitazone in nondiabetic subjects with nonalcoholic steatohepatitis. *Gastroenterology* 2008;135:1176–1184.
101. Cusi K, Orsak B, Bril F, et al. Long-term pioglitazone treatment for patients with nonalcoholic steatohepatitis and prediabetes or type 2 diabetes mellitus: a randomized trial. *Ann Intern Med* 2016;165:305–315.
102. Sanyal AJ, Chalasani N, Kowdley KV, et al. Pioglitazone, vitamin E, or placebo for nonalcoholic steatohepatitis. *N Engl J Med* 2010;362:1675–1685.
103. Kawai M, Rosen CJ. PPAR γ : a circadian transcription factor in adipogenesis and osteogenesis. *Nat Rev Endocrinol* 2010;6:629–636.
104. Inoue I, Shinoda Y, Ikeda M, et al. CLOCK/BMAL1 is involved in lipid metabolism via transactivation of the peroxisome proliferator-activated receptor (PPAR)

- response element. *J Atheroscler Thromb* 2005;12:169–174.
105. Schmutz I, Ripperger JA, Baeriswyl-Aebischer S, et al. The mammalian clock component PERIOD2 coordinates circadian output by interaction with nuclear receptors. *Genes Dev* 2010;24:345–357.
 106. Yang Z, Kim H, Ali A, et al. Interaction between stress responses and circadian metabolism in metabolic disease. *Liver Res* 2017;1:156–162.
 107. Chaix A, Zarrinpar A, Panda S. The circadian coordination of cell biology. *J Cell Biol* 2016;215:15–25.
 108. Ron D, Walter P. Signal integration in the endoplasmic reticulum unfolded protein response. *Nat Rev Mol Cell Biol* 2007;8:519–529.
 109. Fu S, Watkins SM, Hotamisligil GS. The role of endoplasmic reticulum in hepatic lipid homeostasis and stress signaling. *Cell Metab* 2012;15:623–634.
 110. Hetz C, Chevet E, Harding HP. Targeting the unfolded protein response in disease. *Nat Rev Drug Discov* 2013;12:703–719.
 111. Dara L, Ji C, Kaplowitz N. The contribution of endoplasmic reticulum stress to liver diseases. *Hepatology* 2011;53:1752–1763.
 112. Cretenet G, Le Clech M, Gachon F. Circadian clock-coordinated 12 hr period rhythmic activation of the IRE1 α pathway controls lipid metabolism in mouse liver. *Cell Metab* 2010;11:47–57.
 113. Zhang K, Shen X, Wu J, et al. Endoplasmic reticulum stress activates cleavage of CREBH to induce a systemic inflammatory response. *Cell* 2006;124:587–599.
 114. Zhang C, Wang G, Zheng Z, et al. Endoplasmic reticulum-tethered transcription factor cAMP responsive element-binding protein, hepatocyte specific, regulates hepatic lipogenesis, fatty acid oxidation, and lipolysis upon metabolic stress in mice. *Hepatology* 2012;55:1070–1082.
 115. Nakagawa Y, Shimano H. CREBH regulates systemic glucose and lipid metabolism. *Int J Mol Sci* 2018;19(5):1396.
 116. Kim H, Zheng Z, Walker PD, et al. CREBH maintains circadian glucose homeostasis by regulating hepatic glycogenolysis and gluconeogenesis. *Mol Cell Biol* 2017;37(14). e00048–17.
 117. Zheng Z, Kim H, Qiu Y, et al. CREBH couples circadian clock with hepatic lipid metabolism. *Diabetes* 2016;65:3369–3383.
 118. Park JG, Xu X, Cho S, et al. CREBH-FGF21 axis improves hepatic steatosis by suppressing adipose tissue lipolysis. *Sci Rep* 2016;6:27938.
 119. Nakagawa Y, Oikawa F, Mizuno S, et al. Hyperlipidemia and hepatitis in liver-specific CREB3L3 knockout mice generated using a one-step CRISPR/Cas9 system. *Sci Rep* 2016;6:27857.
 120. Nakagawa Y, Satoh A, Tezuka H, et al. CREB3L3 controls fatty acid oxidation and ketogenesis in synergy with PPAR α . *Sci Rep* 2016;6:39182.
 121. Stokkan KA, Yamazaki S, Tei H, et al. Entrainment of the circadian clock in the liver by feeding. *Science* 2001;291(5503):490–493.
 122. Greenwell BJ, Trott AJ, Beytebiere JR, et al. Rhythmic food intake drives rhythmic gene expression more potently than the hepatic circadian clock in mice. *Cell Rep* 2019;27:649–657.
 123. Tripathi A, Debelius J, Brenner DA, et al. The gut-liver axis and the intersection with the microbiome. *Nat Rev Gastroenterol Hepatol* 2018;15:397–411.
 124. Leung C, Rivera L, Furness JB, et al. The role of the gut microbiota in NAFLD. *Nat Rev Gastroenterol Hepatol* 2016;13:412–425.
 125. Frazier K, Chang EB. Intersection of the gut microbiome and circadian rhythms in metabolism. *Trends Endocrinol Metab* 2020;31:25–36.
 126. Thaiss CA, Zeevi D, Levy M, et al. Transkingdom control of microbiota diurnal oscillations promotes metabolic homeostasis. *Cell* 2014;159:514–529.
 127. Thaiss CA, Levy M, Korem T, et al. Microbiota diurnal rhythmicity programs host transcriptome oscillations. *Cell* 2016;167:1495–1510.
 128. Liang X, Bushman FD, FitzGerald GA. Rhythmicity of the intestinal microbiota is regulated by gender and the host circadian clock. *Proc Natl Acad Sci U S A* 2015;112:10479–10484.
 129. Leone V, Gibbons SM, Martinez K, et al. Effects of diurnal variation of gut microbes and high-fat feeding on host circadian clock function and metabolism. *Cell Host Microbe* 2015;17:681–689.
 130. Zarrinpar A, Chaix A, Yooseph S, et al. Diet and feeding pattern affect the diurnal dynamics of the gut microbiome. *Cell Metab* 2014;20:1006–1017.
 131. Uhr GT, Dohnalova L, Thaiss CA. The dimension of time in host-microbiome interactions. *mSystems* 2019;4:e00216–e00218.
 132. Takayasu L, Suda W, Takanashi K, et al. Circadian oscillations of microbial and functional composition in the human salivary microbiome. *DNA Res* 2017;24:261–270.
 133. Collado MC, Engen PA, Bandin C, et al. Timing of food intake impacts daily rhythms of human salivary microbiota: a randomized, crossover study. *FASEB J* 2018;32:2060–2072.
 134. Kaczmarek JL, Musaad SM, Holscher HD. Time of day and eating behaviors are associated with the composition and function of the human gastrointestinal microbiota. *Am J Clin Nutr* 2017;106:1220–1231.
 135. Skarke C, Lahens NF, Rhoades SD, et al. A pilot characterization of the human chronobiome. *Sci Rep* 2017;7(1):17141.
 136. Mukherji A, Kobiita A, Ye T, et al. Homeostasis in intestinal epithelium is orchestrated by the circadian clock and microbiota cues transduced by TLRs. *Cell* 2013;153:812–827.
 137. Zarrinpar A, Chaix A, Xu ZZ, et al. Antibiotic-induced microbiome depletion alters metabolic homeostasis by affecting gut signaling and colonic metabolism. *Nat Commun* 2018;9:2872.
 138. Oh HYP, Ellero-Simatos S, Manickam R, et al. Depletion of gram-positive bacteria impacts hepatic biological functions during the light phase. *Int J Mol Sci* 2019;20(4):812.

139. Wang Y, Kuang Z, Yu X, et al. The intestinal microbiota regulates body composition through NFIL3 and the circadian clock. *Science* 2017;357(6354):912–916.
140. Lavery DJ, Schibler U. Circadian transcription of the cholesterol 7 α hydroxylase gene may involve the liver-enriched bZIP protein DBP. *Genes Dev* 1993;7:1871–1884.
141. Govindarajan K, MacSharry J, Casey PG, et al. Unconjugated bile acids influence expression of circadian genes: a potential mechanism for microbe-host cross-talk. *PLoS One* 2016;11(12):e0167319.
142. Joyce SA, MacSharry J, Casey PG, et al. Regulation of host weight gain and lipid metabolism by bacterial bile acid modification in the gut. *Proc Natl Acad Sci U S A* 2014;111:7421–7426.
143. Segers A, Desmet L, Thijs T, et al. The circadian clock regulates the diurnal levels of microbial short-chain fatty acids and their rhythmic effects on colon contractility in mice. *Acta Physiol (Oxf)* 2019;225(3):e13193.
144. Kohsaka A, Laposky AD, Ramsey KM, et al. High-fat diet disrupts behavioral and molecular circadian rhythms in mice. *Cell Metab* 2007;6:414–421.
145. Ando H, Kumazaki M, Motosugi Y, et al. Impairment of peripheral circadian clocks precedes metabolic abnormalities in ob/ob mice. *Endocrinology* 2011;152:1347–1354.
146. Ho A, Chin A. Circadian feeding and drinking patterns of genetically obese mice fed solid chow diet. *Physiol Behav* 1988;43:651–656.
147. Sookoian S, Castano G, Gemma C, et al. Common genetic variations in *CLOCK* transcription factor are associated with nonalcoholic fatty liver disease. *World J Gastroenterol* 2007;13:4242–4248.
148. Sookoian S, Gemma C, Gianotti TF, et al. Genetic variants of *Clock* transcription factor are associated with individual susceptibility to obesity. *Am J Clin Nutr* 2008;87:1606–1615.
149. Willer CJ, Bonnycastle LL, Conneely KN, et al. Screening of 134 single nucleotide polymorphisms (SNPs) previously associated with type 2 diabetes replicates association with 12 SNPs in nine genes. *Diabetes* 2007;56:256–264.
150. Potter GD, Skene DJ, Arendt J, et al. Circadian rhythm and sleep disruption: causes, metabolic consequences, and countermeasures. *Endocr Rev* 2016;37:584–608.
151. Hatori M, Vollmers C, Zarrinpar A, et al. Time-restricted feeding without reducing caloric intake prevents metabolic diseases in mice fed a high-fat diet. *Cell Metab* 2012;15:848–860.
152. Longo VD, Panda S. Fasting, circadian rhythms, and time-restricted feeding in healthy lifespan. *Cell Metab* 2016;23:1048–1059.
153. Manoogian ENC, Panda S. Circadian rhythms, time-restricted feeding, and healthy aging. *Ageing Res Rev* 2017;39:59–67.
154. Melkani GC, Panda S. Time-restricted feeding for prevention and treatment of cardiometabolic disorders. *J Physiol* 2017;595:3691–3700.
155. Chaix A, Manoogian ENC, Melkani GC, et al. Time-restricted eating to prevent and manage chronic metabolic diseases. *Annu Rev Nutr* 2019;39:291–315.
156. Chaix A, Zarrinpar A, Miu P, et al. Time-restricted feeding is a preventative and therapeutic intervention against diverse nutritional challenges. *Cell Metab* 2014;20:991–1005.
157. Sherman H, Genzer Y, Cohen R, et al. Timed high-fat diet resets circadian metabolism and prevents obesity. *FASEB J* 2012;26:3493–3502.
158. Gill S, Le HD, Melkani GC, Panda S. Time-restricted feeding attenuates age-related cardiac decline in *Drosophila*. *Science* 2015;347(6227):1265–1269.
159. Villanueva JE, Livelo C, Trujillo AS, et al. Time-restricted feeding restores muscle function in *Drosophila* models of obesity and circadian-rhythm disruption. *Nat Commun* 2019;10:2700.
160. Gill S, Panda S. A smartphone app reveals erratic diurnal eating patterns in humans that can be modulated for health benefits. *Cell Metab* 2015;22:789–798.
161. Carlson O, Martin B, Stote KS, et al. Impact of reduced meal frequency without caloric restriction on glucose regulation in healthy, normal-weight middle-aged men and women. *Metabolism* 2007;56:1729–1734.
162. Moro T, Tinsley G, Bianco A, et al. Effects of eight weeks of time-restricted feeding (16/8) on basal metabolism, maximal strength, body composition, inflammation, and cardiovascular risk factors in resistance-trained males. *J Transl Med* 2016;14:290.
163. Sutton EF, Beyl R, Early KS, et al. Early time-restricted feeding improves insulin sensitivity, blood pressure, and oxidative stress even without weight loss in men with prediabetes. *Cell Metab* 2018;27:1212–1221.
164. Hutchison AT, Regmi P, Manoogian ENC, et al. Time-restricted feeding improves glucose tolerance in men at risk for type 2 diabetes: a randomized crossover trial. *Obesity (Silver Spring)* 2019;27:724–732.
165. Haas JT, Francque S, Staels B. Pathophysiology and mechanisms of nonalcoholic fatty liver disease. *Annu Rev Physiol* 2016;78:181–205.
166. Harrison SA, Rinella ME, Abdelmalek MF, et al. NGM282 for treatment of non-alcoholic steatohepatitis: a multicentre, randomised, double-blind, placebo-controlled, phase 2 trial. *Lancet* 2018;391(10126):1174–1185.
167. Hartmann P, Hochrath K, Horvath A, et al. Modulation of the intestinal bile acid/farnesoid X receptor/fibroblast growth factor 15 axis improves alcoholic liver disease in mice. *Hepatology* 2018;67:2150–2166.
168. Feng D, Lazar MA. Clocks, metabolism, and the epigenome. *Mol Cell* 2012;47:158–167.
169. Hirota T, Lee JW, St John PC, et al. Identification of small molecule activators of cryptochrome. *Science* 2012;337(6098):1094–1097.
170. Chen Z, Yoo SH, Takahashi JS. Development and therapeutic potential of small-molecule modulators of circadian systems. *Annu Rev Pharmacol Toxicol* 2018;58:231–252.
171. Lee YS, Cha BY, Saito K, et al. Nobiletin improves hyperglycemia and insulin resistance in obese diabetic ob/ob mice. *Biochem Pharmacol* 2010;79:1674–1683.
172. He B, Nohara K, Park N, et al. The small molecule nobiletin targets the molecular oscillator to enhance

- circadian rhythms and protect against metabolic syndrome. *Cell Metab* 2016;23:610–621.
173. Solt LA, Wang Y, Banerjee S, et al. Regulation of circadian behaviour and metabolism by synthetic REV-ERB agonists. *Nature* 2012;485(7396):62–68.
 174. Chen Z, Yoo SH, Takahashi JS. Small molecule modifiers of circadian clocks. *Cell Mol Life Sci* 2013;70:2985–2998.
 175. Erkekoglu P, Baydar T. Chronopharmacokinetics of drugs in toxicological aspects: a short review for pharmacy practitioners. *J Res Pharm Pract* 2012;1:3–9.
 176. Dallmann R, Brown SA, Gachon F. Chronopharmacology: new insights and therapeutic implications. *Annu Rev Pharmacol Toxicol* 2014;54:339–361.
 177. Bugge A, Feng D, Everett LJ, et al. Rev-erb α and Rev-erb β coordinately protect the circadian clock and normal metabolic function. *Genes Dev* 2012;26:657–667.
 178. Delezie J, Dumont S, Dardente H, et al. The nuclear receptor REV-ERB α is required for the daily balance of carbohydrate and lipid metabolism. *FASEB J* 2012;26:3321–3335.
 179. Adamovich Y, Rousso-Noori L, Zwihaft Z, et al. Circadian clocks and feeding time regulate the oscillations and levels of hepatic triglycerides. *Cell Metab* 2014;19:319–330.
 180. Barclay JL, Shostak A, Leliavski A, et al. High-fat diet-induced hyperinsulinemia and tissue-specific insulin resistance in *Cry*-deficient mice. *Am J Physiol Endocrinol Metab* 2013;304(10):E1053–E1063.
 181. Shimba S, Ogawa T, Hitosugi S, et al. Deficient of a clock gene, brain and muscle Arnt-like protein-1 (BMAL1), induces dyslipidemia and ectopic fat formation. *PLoS One* 2011;6(9):e25231.
 182. Kudo T, Tamagawa T, Kawashima M, et al. Attenuating effect of *Clock* mutation on triglyceride contents in the ICR mouse liver under a high-fat diet. *J Biol Rhythms* 2007;22:312–323.
 183. Chen P, Han Z, Yang P, et al. Loss of clock gene *mPer2* promotes liver fibrosis induced by carbon tetrachloride. *Hepatol Res* 2010;40:1117–1127.
 184. Feng D, Liu T, Sun Z, et al. A circadian rhythm orchestrated by histone deacetylase 3 controls hepatic lipid metabolism. *Science* 2011;331(6022):1315–1319.
 185. Spiegel K, Leproult R, Van Cauter E. Impact of sleep debt on metabolic and endocrine function. *Lancet* 1999;354:1435–1439.
 186. Wang F, Zhang L, Zhang Y, et al. Meta-analysis on night shift work and risk of metabolic syndrome. *Obes Rev* 2014;15:709–720.
 187. Vyas MV, Garg AX, Iansavichus AV, et al. Shift work and vascular events: systematic review and meta-analysis. *BMJ* 2012;345:e4800.
 188. Karlsson B, Knutsson A, Lindahl B. Is there an association between shift work and having a metabolic syndrome? Results from a population based study of 27 485 people. *Occup Environ Med* 2001;58:747–752.
 189. Wilkinson MJ, Manoogian ENC, Zadourian A, et al. Ten-hour time-restricted eating reduces weight, blood pressure, and atherogenic lipids in patients with metabolic syndrome. *Cell Metab* 2020;31:92–104.
 190. Hsieh SD, Muto T, Murase T, et al. Association of short sleep duration with obesity, diabetes, fatty liver and behavioral factors in Japanese men. *Intern Med* 2011;50:2499–2502.
 191. Kim CW, Yun KE, Jung HS, et al. Sleep duration and quality in relation to non-alcoholic fatty liver disease in middle-aged workers and their spouses. *J Hepatol* 2013;59:351–357.
 192. Imaizumi H, Takahashi A, Tanji N, et al. The association between sleep duration and non-alcoholic fatty liver disease among Japanese men and women. *Obes Facts* 2015;8:234–242.
 193. Bernsmeier C, Weisskopf DM, Pflueger MO, et al. Sleep disruption and daytime sleepiness correlating with disease severity and insulin resistance in non-alcoholic fatty liver disease: a comparison with healthy controls. *PLoS One* 2015;10(11):e0143293.
 194. Marin-Alejandro BA, Abete I, Cantero I, et al. Association between sleep disturbances and liver status in obese subjects with nonalcoholic fatty liver disease: a comparison with healthy controls. *Nutrients* 2019;11(2):322.
 195. Liu C, Zhong R, Lou J, et al. Nighttime sleep duration and risk of nonalcoholic fatty liver disease: the Dongfeng-Tongji prospective study. *Ann Med* 2016;48:468–476.
 196. Balakrishnan M, El-Serag HB, Kanwal F, et al. Shiftwork is not associated with increased risk of NAFLD: findings from the National Health and Nutrition Examination Survey. *Dig Dis Sci* 2017;62:526–533.
 197. Katsagoni CN, Papatheodoridis GV, Ioannidou P, et al. Improvements in clinical characteristics of patients with non-alcoholic fatty liver disease, after an intervention based on the Mediterranean lifestyle: a randomised controlled clinical trial. *Br J Nutr* 2018;120:164–175.
 198. Katsagoni CN, Georgoulis M, Papatheodoridis GV, et al. Associations between lifestyle characteristics and the presence of nonalcoholic fatty liver disease: a case-control study. *Metab Syndr Relat Disord* 2017;15:72–79.
 199. Musso G, Cassader M, Olivetti C, et al. Association of obstructive sleep apnoea with the presence and severity of non-alcoholic fatty liver disease. A systematic review and meta-analysis. *Obes Rev* 2013;14:417–431.
 200. Pamidi S, Tasali E. Obstructive sleep apnea and type 2 diabetes: is there a link? *Front Neurol* 2012;3:126.

Received September 8, 2019. Accepted January 29, 2020.

Correspondence


Address correspondence to: Amir Zarrinpar, MD, PhD, 9500 Gilman Dr, MC 0983, University of California, San Diego, La Jolla, California 92093-0983. e-mail: azarrinpar@ucsd.edu; fax: 858-657-5022.

Conflicts of interest

Dr Zarrinpar is a co-founder and equity holder of Tortuga Biosciences. The remaining authors disclose no conflicts.

Funding

Amir Zarrinpar is supported by K08 DK102902, R03 DK114536, R21 MH117780, and R01 HL148801. All authors receive institutional support from P30 DK120515, P30 DK063491, and UL1 TR001442.



Supplementary Figure 1. The gut microbiome and peripheral circadian rhythms. The gut microbiome can modulate peripheral circadian rhythms through several known mechanisms. It can do so through secondary metabolites, such as SCFAs and bile acids, and/or directly through TLRs. DCA, deoxycholic acid; HDCA, hyodeoxycholic acid; HCA, hyocholic acid; LCA, lithocholic acid; MCA, muricholic acid; TCDCA, taurochenodeoxycholic acid; TCA, taurocholic acid; UDCA, ursodeoxycholic acid.